

## **POSTOPERATIVE INSTRUCTIONS**

### **PATELLA OPEN REDUCTION INTERNAL FIXATION**

#### **DIET**

- Begin with clear liquids and light foods (jello, soups, etc.)
- Progress to your normal diet if you are not nauseated

#### **WOUND CARE**

- Maintain your operative dressing. Loosen bandage if swelling of the foot or ankle occurs.
- It is normal for the knee to bleed and swell following surgery – if blood soaks onto the bandage, do not become alarmed – reinforce with additional dressing.
- Remove surgical dressing on the third post-operative day – if minimal drainage is present, apply waterproof band-aids over incisions and change daily.
- Please keep steri-strips in place.
- To avoid infection, keep surgical incisions clean and dry – you may shower by placing waterproof bandages over incision areas. Please remember to change bandages daily.
- NO immersion of operative leg (i.e. bath) until 4 weeks after surgery.
- Brace may come off to shower.

#### **MEDICATIONS**

- Pain medication is injected into the wound and knee joint during surgery – this will wear off within 8-12 hours.
- Colace (Docusate Sodium)
  - This medication is to help with constipation, a common side effect after taking narcotic pain medications (like Norco) and general anesthesia.
  - Take 1 pill in the morning and 1 in the evening to prevent constipation.
  - It is normal to take several days to make a bowel movement after surgery
  - Drink plenty of clear liquids as the anesthesia can cause dehydration/constipation as well.
- Hydrocodone/Acetaminophen (Norco)
  - This is a narcotic medication for pain.
  - This medication is to be taken only AS NEEDED.
  - Plan to stay on a scheduled dose of 1-2 tablets every 4-6 hours for the first 2-3 days.
  - After 2-3 days you should be able to space out or discontinue the medication and transition to Acetaminophen (Tylenol).
  - DO NOT exceed 4,000 mg of Acetaminophen in a 24-hour period.
  - DO NOT drive, drink alcohol, or take Acetaminophen (Tylenol) WHILE taking this.
- Zofran (Ondansetron)
  - This is an anti-nausea medication.
  - It is a dissolving tablet – place it on your tongue, allow it to dissolve, and swallow.
  - Take this as needed every 4-6 hours for the first 2 days after surgery.

Tristan Juhan, MD  
Orthopedic Surgeon  
Office: 408-293-7767  
Email: [tjuhan@goldenstateortho.com](mailto:tjuhan@goldenstateortho.com)  
Website: [tristanjuhanMD.com](http://tristanjuhanMD.com)



- Ibuprofen (i.e. Advil/Motrin) – over the counter
  - As long as you have no personal history of adverse response to anti-inflammatories, use an over-the counter anti-inflammatory such as **Ibuprofen (i.e. Advil/Motrin) 600-800 mg as frequently as every 8 hours** with food to help swelling and pain in addition to the prescribed pain medication.

### **ACTIVITY**

- Non-weightbearing for the first two weeks. Use crutches to assist with walking.
- Elevate the operative leg to chest level whenever possible to decrease swelling.
- Place pillows under knees (i.e., do not maintain knee in a flexed or bent position), but rather place pillows under foot/ankle to elevate leg.
- Do not engage in activities which increase knee pain/swelling (prolonged periods of standing or walking) over the first 7-10 days following surgery.
- Avoid long periods of sitting (without leg elevated) or long distance traveling for 2 weeks.
- May return to sedentary work or school 3-4 days after surgery, if pain is tolerable.

### **ICE THERAPY**

- Begin icing immediately after surgery.
- Use icing machine for up to 20 minutes out of every hour until your first post-op visit.
- If you do not purchase an ice machine, please use ice packs to ice the surgical site. Ice packs can be applied for up to 20 minutes out of every hour until your first post-op visit.
- If braced, loosen brace to avoid added pressure.

### **EXERCISES**

- Formal physical therapy (PT) will begin within a week of your surgery. A prescription will be given to you. Please give the therapist the REHABILITATION form from our office, also located on my website.
- Begin exercises 24 hours after surgery (straight leg raises, quad sets, and ankle pumps) unless otherwise instructed.
- Complete exercises 3-4 times daily until your first post-op visit
- Do ankle pumps continuously throughout the day to reduce the possibility of a blood clot in your calf (extremely uncommon).
- NO KNEE MOTION AT THIS TIME

### **FOLLOW-UP CARE/QUESTIONS**

- Dr. Juhan will call you (typically on your first day after surgery) to address any questions or concerns. If you have not been contacted within 48 hours of surgery, please call the office at **408-293-7767**
- If you do not already have a post-operative appointment scheduled, please contact the office during normal office hours and ask for appointment scheduling at **408-293-7767**.
- For any other questions or concerns please contact Dr. Juhan via email ([tjuhan@goldenstateortho.com](mailto:tjuhan@goldenstateortho.com)) at the office (**408-293-7767**)

Tristan Juhan, MD  
Orthopedic Surgeon  
Office: 408-293-7767  
Email: [tjuhan@goldenstateortho.com](mailto:tjuhan@goldenstateortho.com)  
Website: [tristanjuhanMD.com](http://tristanjuhanMD.com)



**\*\*EMERGENCIES\*\***

Contact Dr. Juhan at the office (**408-293-7767**) or by email ([tjuhan@goldenstateortho.com](mailto:tjuhan@goldenstateortho.com)) if any of the following are present:

- Painful swelling or numbness
- Unrelenting pain
- Fever (note – it is normal to have a low-grade fever (101° and under) for the first day or two following surgery) or chills
- Redness around incisions
- Color change in wrist or hand
- Continuous drainage or bleeding from incision (a small amount of drainage is expected)
- Difficulty breathing
- Excessive nausea/vomiting

**Proceed to the nearest emergency room if you have an emergency that requires immediate attention**